

AD 741394

AD _____


4

PARASITIC NEMATODES OF SOUTHEAST ASIA
AS POTENTIAL ZOONOSES

FINAL REPORT

GERALD D. SCHMIDT

MARCH, 1972



Supported by

U. S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Washington, D. C. 20315

Contract No. DADA17-68-C-8094
University of Northern Colorado
Greeley, Colorado 80631

This document has been approved for public release and sale;
its distribution is unlimited.

The findings in this report are not to be construed as an
Official Department of the Army position unless so designa-
ted by other authorized documents.

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22161

No Record of #3
#2 - AD 741394

9

PARASITIC NEMATODES OF SOUTHEAST ASIA
AS POTENTIAL ZOONOSES

FINAL REPORT

CEPALD D. SCHMIDT

MARCH, 1972

Supported by

U. S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Washington, D. C. 20315

Contract No. DADA17-68-C-8094
University of Northern Colorado
Greeley, Colorado 80631

This document has been approved for public release and sale;
its distribution is unlimited.

The findings in this report are not to be construed as an
Official Department of the Army position unless so designated
by other authorized documents.

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified.)

1. ORIGINATING ACTIVITY (Corporate author)	2a. REPORT SECURITY CLASSIFICATION
University of Northern Colorado Greeley, Colorado 80631	2b. GROUP

3. REPORT TITLE

PARASITIC NEMATODES OF SOUTHEAST ASIA AS POTENTIAL ZOONOSES

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)
Final - 1 March, 1968 - 27 March, 1972

5. AUTHOR(S) (First name, middle initial, last name)

Gerald D. Schmidt

6. REPORT DATE 1 March, 1972	7a. TOTAL NO. OF PAGES Eight	7b. NO. OF REFS Twenty-two
8. CONTRACT OR GRANT NO. DADA17-68-C-8094	8a. ORIGINATOR'S REPORT NUMBER(S) Four	
9. PROJECT NO.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.		
d.		
10. DISTRIBUTION STATEMENT DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY U. S. Army Medical Research and Development Command. Surgeon General Washington, D. C. 20315	

13. ABSTRACT

Studies of the parasitic nematodes of Southeast Asia collected by NAMRU-2, have been completed. Twenty publications have resulted from this research. Host-parasite and locality records have been published, as have descriptions of numerous poorly-known species. In addition, 37 species of nematodes new to science has been described and named. Six new genera have been established. Recommendations for continued support of this type of faunal exploration are presented.

Security Classification

KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Nematodes						
Parasites						
Zoonoses						
Palawan, nematodes of						
Borneo, nematodes of						
Taiwan, nematodes of						

Summary

Studies of the parasitic nematodes of Southeast Asia collected by NAMRU-2 have been completed. Twenty publications have resulted from this research. Host-parasite and locality records have been published, as have descriptions of numerous poorly-known species. In addition, 37 species of nematodes new to science have been described and named. Six new genera have been established. Recommendations for continued support of this type of faunal exploration are presented.

Statement of the Problem

This study was undertaken as a part of the Southeast Asia faunal exploration undertaken by U. S. Navy Medical Research Unit No. 2., Taipei, Taiwan. This epidemiological team collected thousands of vertebrates and invertebrates in efforts to determine the ecological and zoological aspects of endemic and epidemic diseases. The primary problem to be solved in such investigations is the taxonomic one, involving descriptions of the organisms in such a manner that they may be readily recognized by future workers. To this end, the parasitic helminths collected by NAMRU-2 during field operations on Taiwan and during expeditions to Borneo, Philippines, Korea, Pakistan and Solomon Islands have been entrusted to specialists for study. Most parasitic nematodes were sent to the present investigator, who was awarded a contract, DADA17-68-6-8094, by the U. S. Army Medical Research and Development Command, to help defray the expenses of the study.

This constitutes the final report on the research activities supported by this contract.

Approach to the Problem

Approximately one thousand vials of preserved nematodes, collected from fishes, amphibians, reptiles, birds and mammals were studied by conventional parasitological techniques. Species new to science were described and named. Poorly known species were redescribed, often with revision of the higher categories of classification and with identification keys to the genera and species. New host and distribution records were recorded, and those parasites with obvious potential for infecting man were noted. The results were published in a variety of professional journals, coauthored with Captain Robert E. Kuntz, of NAMRU-2.

Results

The results of these studies have been published in the series "Nematode Parasites of Oceanica", parts 1-XX. (The first four parts were published prior to the Army contract). These publications are listed below under Selected Bibliography, and should be consulted for detailed results.

The following new taxa were described in these papers:

New Subfamily

Arthrocephelinae Schmidt et Kuntz, 1968

New Genera

Calypsostrongylus Schmidt et Kuntz, 1967
Oceanicucullanus Schmidt et Kuntz, 1969
Oceanifilaria Schmidt et Kuntz, 1970
Madelinema Schmidt et Kuntz, 1971
Cordonema Schmidt et Kuntz (in press).
Smetaleksanema Schmidt et Kuntz, (in press).

New Species

Brevistriata sundasciuri Schmidt, Myers et Kuntz, 1967
Calypsostrongylus ogeni Schmidt, Myers et Kuntz, 1967
Arthrostoma vampira Schmidt et Kuntz, 1968
Syphacia oceanica Schmidt et Kuntz, 1968
Syphacia coli Schmidt et Kuntz, 1968
Syphacia critesi Schmidt et Kuntz, 1969
Oceanicucullanus pacifica Schmidt et Kuntz, 1969
Camallanus marinus Schmidt et Kuntz, 1969
Spinitectus palawanensis Schmidt et Kuntz, 1969
Cucullanus lutjani Schmidt et Kuntz, 1969
Fleleyella confusa Schmidt et Kuntz, 1969
Icosiella hoogstraali Schmidt et Kuntz, 1969
Oceanifilaria verrucosa Schmidt et Kuntz, 1970
Aprocta calliderma Schmidt et Kuntz, 1970
Parornithofilaria scinini Schmidt et Kuntz, 1970
Parornithofilaria hepatica Schmidt et Kuntz, 1970
Capillaria parusi Wakelin, Schmidt et Kuntz, 1970
Capillaris madseni Wakelin, Schmidt et Kuntz, 1970
Capillaria javicensis Wakelin, Schmidt et Kuntz, 1970
Capillaria pitti Wakelin, Schmidt et Kuntz, 1970
Capillaria anthracocerosi Wakelin, Schmidt et Kuntz, 1970
Inglosionema mawsonae Schmidt et Kuntz, 1971
Madelinema angelae Schmidt et Kuntz, 1971
Tetrameres robusta Schmidt et Kuntz, 1971
Acuraria kinsellai Schmidt et Kuntz, 1971
Rusguniella microcordonis Schmidt et Kuntz, 1971
Subulura helicospicula Schmidt et Kuntz, 1971
Ceratospirura inglisi Schmidt et Kuntz, 1971
Paraheterotyphlum ophiophagos Schmidt et Kuntz, (in press)
Caenorhabditis avicola Schmidt et Kuntz, (in press)
Heterakis vexana Schmidt, Inglis et Kuntz, (in press)
Viktorocara acholonui Schmidt et Kuntz, (in press)
Ornithostrongylus vetterlingi Schmidt et Kuntz, (in press)
Cordonema venusta Schmidt et Kuntz, (in press)
Skrjabinoclava raliae Schmidt et Kuntz, (in press)
Skrjabinoclava aurorae Schmidt et Kuntz, (in press)

Discussion and Conclusions

The only species of nematode that is known to be a human pathogen which was found in this study is Gnathostoma spinigerum, which was found encysted in frogs in Palawan. The ingestion of raw frog anywhere in the Orient is to be avoided.

Anasakis-type larvae are extremely abundant in the marine fishes of the Philippines. The type of larva is known to cause gastric tumors whenever marine fish is eaten raw. Raw fish is likewise to be avoided throughout the Orient and oceanic islands.

Capillaria philippensis was not found in this study, nor was Angiostrongylus cantonensis. Both are known to inhabit the areas sampled, however, which shows that this survey, extensive as it was, was still an incomplete sampling of the parasites of the region.

When viewed from the ecological-epidemiological viewpoint, the first task of any zoonosis survey is taxonomic. It would therefore appear that the present study accomplished its mission: the recognition of a substantial number of endemic species of parasitic nematodes.

Recommendations

It is recommended that further sampling of the parasitic fauna of Southeast Asia be accomplished, not only of nematodes, but also of cestodes, Acanthocephala, trematodes and protozoa. Even more importantly, financial support should be available for the specialists who work up the collections made by the Government. It is such cooperation between government and civilian workers that our final goal will be accomplished: global eradication of disease.

Bibliography

1. Schmidt, G. D., B. J. Myers, and R. E. Kuntz. 1967. Nematode parasites of Oceanica. I. Brevistriata sundasciuri sp. n. and Calypsostrongylus ogdeni gen. et sp. n. (Heligmosomatidae: Longistriatinae) from squirrels of Palawan and Taiwan. *J. Parasitol.* 53: 613-617.
2. _____, and R. E. Kuntz. 1967. Nematode parasites of Oceanica. II. Redescription of Rictularia whartoni, 1931, and notes on other species from Palawan, P. I. *J. Parasitol.* 53: 1281-1284.
3. _____, and _____. 1968. Nematode parasites of Oceanica. III. Arthrostoma vampira sp. n., with a reconsideration of those hookworms having articulated buccal capsules. *J. Parasitol.* 54: 372-376.
4. _____, and _____. 1968. Nematode parasites of Oceanica. IV. Oxyurids of mammals of Palawan, P. I., with descriptions of four new species of Syphacina. *Parasitology* 58: 845-854.
5. _____, and _____. 1969. Nematode parasites of Oceanica. V. Four new species from fishes of Palawan, P. I., with a proposal for Oceanicucullanus gen. nov. *Parasitol.* 59: 389-396.
6. _____, and _____. 1970. Nematode parasites of Oceanica. VI. Polevella confusa sp. nov., Icosiella hoogstraali sp. nov. (Filarioidea), and other species from Philippine amphibians. *Parasitol.* 59: 885-889.
7. _____, and _____. 1970. Nematode parasites of Oceanica. VII. New records from wild and domestic chickens (Gallus galus) from Palawan (Philippine Islands), Sabah (Malaysia) and Taiwan. *Avian Diseases* 14: 184-187.
8. _____, and _____. 1970. Nematode Parasites of Oceanica. VIII. Filariids of birds, with a new genus and four new species. *Parasitol.* 60: 313-326.
9. Wakelin, D., G. D. Schmidt, and R. E. Kuntz, 1970. Nematode parasites of Oceanica. IX. Capillariids from passeriform hosts collected in Taiwan. *Parasitology*. 61: 465-474.
10. _____, _____, and _____. 1971. Nematode parasites of Oceanica X. Capillariids from avian hosts collected in Palawan and Borneo. *Parasitology* 62: 1-10.
11. Schmidt, G. D., and R. E. Kuntz. 1971. Nematode parasites of Oceanica. XI. Madelininema arangelae gen. et. sp. n., and Inglisonema mawsonic sp. n. (Heterakoidea: Inglisonematidae) from birds. *J. Parasitol.* 57: 479-484.

12. _____, W. G. Inglis, and R. E. Kuntz. (In press). Nematode parasites of Oceanica. XII. A review of Heterakis species, particularly from birds of Taiwan and Palawan. Rec. S. Aust. Mus.

13. _____, and R. E. Kuntz. 1971. Nematode parasites of Oceanica. XIII. Subuluridae, Thelaziidae and Acuariidae of birds. Parasitology 63: 91-99.

14. _____, and _____. 1971. Nematode parasites of Oceanica XIV. Tetrarmeridae and Heterocheilidae of birds. Trans. Am. Micro. Soc. 90: 325-329.

15. _____, and _____. 1971. Nematode parasites of Oceanica. XV. Acuariidae, Streptocaridae, and Seuratidae of birds. Proc. Helm. Soc. Wash. 38: 217-223.

16. _____, and _____. (In press). Nematode parasites of Oceanica. XVI. Cordonema venusta gen. et. sp. n., and Skriabinoclava spp., (Acuariidae: Echinuriinae), from birds. Parasitology.

17. _____, and _____. (In press). Nematode parasites of Oceanica. XVII. Schistorophidae, Spiruridae, Phsalopteridae and Trichostrongylidae of birds. Parasitology.

18. _____, and _____. (In press). Nematode parasites of Oceanica. XVIII. Gaenorhabditis avicola sp. n. (Rhabditidae) found in a bird from Taiwan. Proc. Helm. Soc. Wash.

19. _____, and _____. 1972. Nematode parasites of Oceanica. XIX. Report on a collection from Philippine reptiles. Trans. Am. Micro. Soc. 91: 63-66.

20. _____, and _____. (In press). Nematode parasites of Oceanica. XX. Paraheterotyphlum ophiophagorum sp. n. (Heterocheilidae), from the banded yellow-lip sea snake, Laticephala colubrina. Trans. Am. Micro. Soc.